

HIGH-DENSITY CA-RICH BRINES IN OH-RICH TOPAZ FROM KYANITE QUARTZITE FROM THE SU-LU UHP METAMORPHIC BELT (EASTERN CHINA).

FERRANDO, S.¹, FREZZOTTI, M.L.², COMPAGNONI, R.³

¹Dipartimento di Scienze Mineralogiche e Petrologiche, Via Valperga Caluso 35, 10125 Torino.

²Dipartimento di Scienze della Terra, Via Laterina 8, 53100 Siena.

³Dipartimento di Scienze Mineralogiche e Petrologiche, Via Valperga Caluso 35, 10125 Torino.

E-mail: simona.ferrando@unito.it

Metamorphic OH-rich topaz has been recently reported from kyanite quartzite at Hushan, belonging to the high-pressure (HP) to ultra-high pressure (UHP) eclogite-facies metamorphic belt of Su-Lu, eastern China (Zhang et al., 2002). The stability of this mineral at quartz- and coesite-facies conditions has been confirmed by experimental works (Wunder et al., 1993; Wunder et al., 1999).

Isopleths for the reactions Al-silicate + fluid = topaz, diaspore + quartz = topaz, and diaspore + quartz = fluid + Al-silicate were calculated thermodynamically (Holland, Powell, 1998; Connolly, 1990) by considering both the hydroxyl contents of the topaz and variable $X_{H_2O} = H_2O/(H_2O + CO_2)$ in the fluid phase. The stability of the F-OH topaz solid solution series results to be strongly controlled by both P-T conditions and fluid composition. In particular, the topaz stability field is reduced by the CO_2 increase in the fluid phase.

The studied Hushan kyanite quartzite consists of quartz, kyanite, paragonite, OH-rich topaz, accessory rutile, pyrite, zircon, apatite, and rare barite. The presence in kyanite of rare polycrystalline quartz aggregates suggests the former occurrence of coesite. The $X_{OH} = OH/(OH+F)$ ratio in topaz is quite homogeneous and averages 0.28 (Alberico et al., 2002). Peak metamorphic conditions are consistent with climax conditions estimated for the associated eclogites ($T = 800 \pm 80$ °C and $P = 3.2 - 4.0$ GPa; Zhang et al., 2000). Microstructural relationships clearly indicate that the OH-rich topaz formed at the expense of kyanite during early decompression by a reaction such as: kyanite + fluid = topaz.

OH-rich topaz crystals contain extremely abundant primary (in the sense of Roedder, 1984) fluid inclusions, having rounded or negative crystal shapes. The inclusions, ranging in size from < 5 to $20 \mu m$, are water dominated with minor CO_2 (5 % in volume) and may contain an anhydrite crystal. Evidence of post-trapping changes, such as partial decrepitation, are often observed. Resetted inclusions contain larger CO_2 bubble ($CO_2 = 10$ to 50% of the inclusion total volume) and in a few rare cases a single chloride crystal.

The aqueous part of the preserved fluid inclusions freezes below $-60^\circ C$. The eutectic temperature cannot be accurately measured, but at around $-45^\circ C$ a liquid phase is always observed, which suggests the presence of Ca^{2+} in the fluid. The temperature of hydrohalite melting (T_{mhy}) occurs between -37.0 and $-32.0^\circ C$ with most data at $-34.4^\circ C$, and ice final melting temperatures (T_{mice}) between -17.0 and -13.0 with most data at $-13.7^\circ C$; CO_2 melts (T_{mCO_2}) instantaneously at $\approx -56.6^\circ C$, and homogenisation, always to the liquid phase (T_{hLCO_2}), is recorded between -5.8 and $9.7^\circ C$. Original trapped fluid composition corresponds to a high-density ($1.14 - 1.16 g/cm^3$) Ca-rich brine containing minor CO_2 ($X_{H_2O} = 0.87$, $X_{CO_2} = 0.04$, $X_{NaCl} = 0.01$; $X_{CaCl_2} = 0.08$); mean salinity for the aqueous part of the fluid is 2.5 wt % NaCl and 15 wt % $CaCl_2$ (Bakker, 1999). Resetted inclusions have considerably lower density ($1.11 - 0.66 g/cm^3$), higher CO_2 content ($X_{CO_2} = 0.06-0.09$) and variable salinity (mean values: 4 wt % NaCl + 11 wt % $CaCl_2$), indicating that some preferential water leakage occurred during decrepitation processes.

The collected data indicate that the fluid phase that assisted the topaz growth at the expense of kyanite was a Ca-dominated medium-salinity brine, containing traces of CO_2 . The isochores, calculated from the fluid inclusions with the highest densities (Bakker, 1999), indicate a minimum pressure of 2.8 GPa at the post-climactic temperature of $800^\circ C$, estimated from the Su-Lu coesite-eclogite (Nakamura, Hirajima, 2000). The preservation of high-pressure fluid inclusions in microdomains within the topaz crystals, further shows this mineral as a good host for the study of fluid inclusions in high-pressure metamorphic rocks.

A petrogenetic grid, calculated with thermodynamic approach (Holland, Powell, 1998; Connolly, 1990) and considering both topaz ($X_{OH} = 0.30$) and fluid phase composition ($X_{H_2O} = 0.90$), indicates that the OH-rich topaz grew during the early decompression, at the coesite to quartz transition, on the P-T path estimated for the Su-Lu UHPM rocks (Nakamura, Hirajima, 2000).

References

- ALBERICO, A., FERRANDO, S., IVALDI, G., FERRARIS, G. (2002): X-ray single-crystal structure refinement of an OH-rich topaz from Sulu UHP terrane (Eastern China). Structural foundation of the correlation between cell parameters and fluorine content. Submitted to European Journal of Mineralogy.
- BAKKER, R. J. (1999) Optimal Interpretation of microthermometrical data from fluid inclusions: thermodynamic modelling and computer programming. Habilitation, Ruprecht-Karls-University Heidelberg (Germany), 1-50.
- CONNOLLY, J. A. D. (1990): Multivariable phase diagrams: an algorithm based on generalized thermodynamics. *Am. J. Sci.*, **290**, 666-718.
- HOLLAND, T. J. B., POWELL, R. (1998): An internally consistent thermodynamic dataset for phases of petrological interest. *J. Metam. Geol.*, **16**, 309-343.
- NAKAMURA, D., HIRAJIMA, T. (2000): Granulite-facies overprinting of ultrahigh-pressure metamorphic rocks, Northeastern Su-Lu Region, Eastern China. *Journal of Petrology*, **41/4**, 563-582.
- ROEDDER E. (1984): Fluid inclusions. P. Ribbe (ed): *Reviews in Mineralogy*, Volume 12. Mineralogical Society of America, Washington, D. C., 646 pp.
- WUNDER, B., ANDRUT, M., WIRTH, R. (1999): High-pressure synthesis and properties of OH-rich topaz. *Eur. J. Mineral.*, **11**, 803-813.
- WUNDER, B., RUBIE, D. C., ROSS II, C. R., MEDENBACH, O., SEIFERT, F., SCHREYER, W. (1993): Synthesis, stability, and properties of $\text{Al}_2\text{SiO}_4(\text{OH})_2$: a fully hydrated analogue of topaz. *Amer. Mineral.*, **78**, 285-297.
- ZHANG, R. Y., LIOU, J. G., SHU, J. F. (2002): Hydroxyl-rich topaz in high-pressure and ultrahigh-pressure kyanite quartzites, with retrograde woodhouseite, from the Sulu terrane, eastern China. *Amer. Mineral.*, **87**, 445-453.
- ZHANG, Z., XU, Z., XU, H. (2000): Petrology of ultrahigh-pressure eclogites from the ZK703 drillhole in the Donghai, eastern China. *Lithos*, **52**, 35-50.